

UNCLASSIFIED

AD 296 581

*Reproduced
by the*

**ARMED SERVICES TECHNICAL INFORMATION AGENCY
ARLINGTON HALL STATION
ARLINGTON 12, VIRGINIA**



UNCLASSIFIED

NOTICE: When government or other drawings, specifications or other data are used for any purpose other than in connection with a definitely related government procurement operation, the U. S. Government thereby incurs no responsibility, nor any obligation whatsoever; and the fact that the Government may have formulated, furnished, or in any way supplied the said drawings, specifications, or other data is not to be regarded by implication or otherwise as in any manner licensing the holder or any other person or corporation, or conveying any rights or permission to manufacture, use or sell any patented invention that may in any way be related thereto.

63-2-4

USNRDL-TR-605

Copy 112
3 January 1963

296 581

296 581

MOUSE FETAL LIVER: A SOURCE OF
IMMUNOLOGICALLY REACTIVE CELLS

by
M.L. Tyan
L.J. Cole

CATALOGED BY ASTIA
AS AD NO.

ASTIA
RECEIVED
FEB 25 1963
RECEIVED
TISA

U.S. NAVAL RADIOLOGICAL
DEFENSE LABORATORY
SAN FRANCISCO 24, CALIFORNIA

12ND. P7463

EXPERIMENTAL PATHOLOGY BRANCH
L.J. Cole, Head

BIOLOGICAL AND MEDICAL SCIENCES DIVISION
E.L. Alpen, Head

ADMINISTRATIVE INFORMATION

This work was accomplished under the Bureau of Medicine and Surgery Task MR005.08-5200, Subtask 3, Technical Objective AW-6, as described in the U.S. Naval Radiological Defense Laboratory Annual Report to the Bureau of Medicine and Surgery (OPNAV Form 3910-1) of 31 December 1962, and is listed in the U.S. Naval Radiological Defense Laboratory Technical Program Summary for Fiscal Years 1963-1965 of 1 November 1962 under Program A3, Problem 2, entitled "Nuclear Warfare Aspects of Whole Body Ionizing Radiation." This study was supported through funds provided by the Bureau of Medicine and Surgery, and the Defense Atomic Support Agency under NWER Program A 4c, Subtask 03.027.

Eugene P. Cooper
Eugene P. Cooper
Scientific Director

E. B. Roth
E. B. Roth, CAPT USN
Commanding Officer and Director

ABSTRACT

Male and female AHeJ mice were mated and the resulting pregnancies were surgically interrupted at various stages of gestation. The embryos were dissected and identical organs (liver, thymus, gut, placenta, and umbilicus) from each separate pregnancy were pooled and injected i. p. into sublethally irradiated (500 rad) F_1 hybrids (either BALB/c x A) F_1 or (C57L x A) F_1). Death of the F_1 hybrids within 60 days was the criterion for the presence of immunologically competent cells in the inoculum. At the end of this period the survivors were sacrificed, their spleens and lymph nodes homogenized, and injected i. p. into sublethally irradiated F_1 hybrids of the other type. Again death within 60 days was the criterion for the presence of immunologically competent cells in the inoculum.

One death occurred among the primary hosts (placenta 1/8). Immunologically competent cells (AHeJ) were detected in secondary recipients of 2nd trimester fetal liver (3/3), 2nd trimester thymus (3/3), and 3rd trimester fetal liver (11/12). No immunologically competent cells (AHeJ) were found in secondary recipients of 3rd trimester and newborn thymus (1/14). Deaths occurred among secondary hosts of fetal gut (3/5), umbilicus (1/3), and placenta (2/9). There were two deaths (2/29) among the secondary control mice. It is concluded that 2nd and

3rd trimester mouse fetal liver, 2nd trimester fetal thymus, and perhaps other fetal tissues contain potential immunologically competent cells. The theoretical significance of these data is discussed.

SUMMARY

The Problem:

Recent observations have focused attention upon the mammalian embryonic and neonatal thymus as a critical organ in the ultimate development of immunological competence. However, other data indicate that mouse fetal liver may be a source of potential immunologically competent cells. The present work adds further evidence for the presence of potential immunologically competent cells in both mouse fetal liver and thymus.

The Findings:

Male and female AHeJ mice were mated and the resulting pregnancies were surgically interrupted at various stages of gestation. The embryos were dissected and identical organs (liver, thymus, gut, placenta, and umbilicus) from each separate pregnancy were pooled and injected i. p. into sublethally irradiated (500 rad) F_1 hybrids (either (BALB/c x A) F_1 or (C57L x A) F_1). Death of the F_1 hybrids within 60 days was the criterion for the presence of immunologically competent cells in the inoculum. At the end of this period the survivors were sacrificed, their spleens and lymph nodes homogenized, and injected i. p. into sublethally irradiated F_1 hybrids of the other type. Again death within

60 days was the criterion for the presence of immunologically competent cells in the inoculum.

One death occurred among the primary hosts (placenta 1/8). Immunologically competent cells (AHeJ) were detected in secondary recipients of 2nd trimester fetal liver (3/3), 2nd trimester thymus (3/3), and 3rd trimester fetal liver (11/12). No immunologically competent cells (AHeJ) were found in secondary recipients of 3rd trimester and newborn thymus (1/14). Deaths occurred among secondary hosts of fetal gut (3/5), umbilicus (1/3), and placenta (2/9). There were two deaths (2/29) among the secondary control mice. It is concluded that 2nd and 3rd trimester mouse fetal liver, 2nd trimester fetal thymus, and perhaps other fetal tissues contain potential immunologically competent cells. The theoretical significance of these data is discussed.

INTRODUCTION

Recent observations have focused attention upon the embryonic and neonatal thymus as a critical organ in the ultimate development of immunological competence. Kalmutz (1) demonstrated that the opossum embryo was able to react against antigenic stimuli as soon as the thymus had matured, but prior to maturation of other "lymphoid organs". Miller (2), Martinez et al (3) and others (4-7) have shown that mice and rats thymectomized within 24 hours of birth have a markedly reduced life span and a broad impairment of their immunological responsiveness. It has been suggested (2-9) that the fetal thymus and its descendent lympho-epithelial cells, either by the centrifugal distribution of thymocytes which give rise directly to functionally active "collateral" cells, or by some as yet unknown process which promotes proliferation and maturation of locally arising lymphocytic cells, govern the immunological reactivity of the mammalian organism.

However, rodents protected against the lethal effects of X-radiation by the intravenous infusion of isogenic or allogeneic fetal liver cells manifest none of the stigmata of immunological incompetence (10-14). Moreover, Doria et al (15), using 12-18 day mouse embryos, demonstrated that fetal liver "contained potential or actual antibody forming cells" which when transplanted into lethally irradiated hosts, conferred upon them a competent immune system of the donor type. The present communication is intended to add further evidence for the presence of potential

immunologically competent cells in both fetal liver and thymus, and to discuss briefly the possible significance of these and the above cited data.

MATERIALS AND METHODS

The presence of potential immunologically reactive cells in various fetal tissues was determined by the parental- F_1 hybrid, graft-versus-host method of Cole (16), with the following modification. It could be assumed that any immature lymphoid cells which might be present in an inoculum of parental strain fetal tissue would rapidly develop tolerance to their sublethally irradiated F_1 hybrid host, and, as a result, their presence would not be detected. If, however, maturation of these potential immunologically competent cells occurred in the F_1 hybrid host, it should be possible to demonstrate them upon transfer to an F_1 hybrid of another type (one parent in common).

Twelve week old mice, either (C57L x A) F_1 or (BALB/c x A) F_1 (called LAF $_1$ and CAF $_1$ respectively), which had received 500 rad whole body X radiation just prior to the i. p. injection of parental strain fetal tissue, were used as primary hosts. Death of these sublethally irradiated F_1 hybrids within 60 days was the criterion for the presence of immunologically competent cells in the inoculum. At the end of this period, each survivor was sacrificed; its spleen and lymph nodes extirpated, lightly homogenized in cold Tyrode's solution, and the cell suspension injected into 1 or 2 sublethally irradiated F_1 hybrids of the other type. Again,

death of the secondary host within 60 days indicated the presence of immunologically reactive parental strain cells in the inoculum. F_1 hybrids (of the same type as the primary host), which had received 500 rad whole body X radiation but no injection of fetal tissue, served as primary radiation controls. Sublethally irradiated F_1 hybrids (of the same type as the secondary host), which received an i. p. injection of spleen and lymph nodes from the primary control animals, were the controls for the secondary hosts. The irradiated mice were housed 4-10 per cage. The diet was Purina Lab Chow, and water (containing 1% Neomycin) was given ad libitum.

Male and female AHeJ mice were mated and the resultant pregnancies were surgically extirpated at various stages of gestation. The embryos were separated from the other products of conception without contamination by maternal blood or tissue, and the various organs under study were dissected free in the cold. Identical organs from a single pregnancy were pooled in cold Tyrode's solution and the cells were dissociated by gentle aspiration through a 23 gauge needle. Where placental tissue was used, the organs were grossly dissected free of maternal decidua, finely minced in cold Tyrode's solution, and the cells were dissociated, by gentle aspiration through a 20 gauge needle. It is doubtful that the placental tissue was entirely free of maternal contamination. The cell suspensions were then injected intraperitoneally into 1 or 2 irradiated F_1 hybrids. Each pregnancy was comprised of from three to nine embryos

of from 11 days gestation to the newly born. The fetal organs used were thymus, liver, gut, umbilicus and placenta. Primary recipients of fetal thymus received from $1 - 10 \times 10^6$ nucleated cells per animal while recipients of fetal liver received from $25 - 150 \times 10^6$ nucleated cells. No attempt was made to quantitate the number of gut, umbilicus or placenta cells given.

RESULTS

As can be seen in Table I, one death occurred among the primary hosts (placenta 1/8), indicating: (1) the absence of mature immunologically competent cells in the fetal tissue injected, (2) inadequate numbers of these cells were given, or (3) the potential lymphoid cells in the inoculum were incapable of reactivity in this system. However, deaths did occur among the secondary recipients of fetal liver and thymus from 11 or 12 day gestations (3/3 and 3/3), and among secondary hosts of third trimester fetal liver (11/12). There was one death among the secondary recipients of third trimester and new born thymus (1/14). The deaths occurring among secondary recipients of fetal gut (3/5), umbilicus (1/3) and placenta (2/9) may be explained by contamination with fetal liver, which occurred occasionally during dissection, in the case of gut and umbilicus, and by contamination with maternal tissue in the case of placenta. These latter findings, however, warrant further investigation. There were two deaths (2/29) among the secondary control animals (both occurring in mice housed in the same cage).

DISCUSSION

On the basis of these data and of the observations of Doria et al (15), it is apparent that fetal liver and 2nd trimester fetal thymus contain potential immunologically competent cells, which under the proper conditions are capable of immunological reactivity. These cells were characterized by an apparent initial absence of immunological reactivity in this system and/or an increased ability to become "tolerant" of allogeneic living cells. It should be noted that cells of similar potential were not found in 3rd trimester thymus or in newborn thymus. It is of interest that Delmaso et al (6) in their study of the immunological competence of spleen cells from mice thymectomized during early neonatal life, observed that these cells manifested no reactivity in primary hosts. The present data suggest that such cells might well have exhibited immunological reactivity, had they been transferred to suitable secondary hosts.

That the thymus is not the only source of cells possessing potential immunological competence in the mouse fetus now seems clear. Mouse fetal liver and, perhaps, other fetal tissues (i.e., gut, spleen, lymph nodes) may contain cells which, under the proper circumstances, and when exposed to appropriate stimuli, are capable of immunological reactivity. The mammalian "immune system" may, therefore, be composed of distinct "lines" of cells, phylogenetically derived at different times, which manifest diversity in degree, mode and specificity of their reactivity,

TABLE I

POTENTIAL IMMUNOLOGICALLY COMPETENT CELLS
IN THE LIVER AND THYMUS OF A/H₂J MOUSE EMBRYOS

TRIMESTER OF PREGNANCY	FETAL CELLS INJECTED (A/H ₂ J)	60-DAY MORTALITY(NO/TOTAL)		MEAN SURVIVAL TIME (days)
		Primary Host (LAF ₁ or CAF ₁)	Secondary Host (LAF ₁ or CAF ₁)	
Second [*]	Thymus	0/2	3/3	15
	Liver	0/2	3/3	26
	Placenta	0/1	1/1	47
	Gut	0/1	0/1	
	None	0/6	0/6	
Third ^{**}	Thymus	0/8	1/12	34
	Liver	0/8	11/12	19
	Placenta	1/8	2/9	16
	Gut	0/5	3/5	16
	Umbilicus	0/3	1/3	10
	None	0/32	2/19	10
Newborn ^Δ	Thymus	0/1	0/2	
	None	0/6	0/4	

^{*}Pooled data from two separate pregnancies (11 and 12 day gestation)

^{**}Pooled data from seven separate pregnancies (18 to 20 day gestation)

^ΔOne litter of seven

but which in the adult animal occur as an apparently homogeneous cell population. It is proposed that during the 3rd trimester of pregnancy in mice, the lympho-epithelial cells of the thymus differentiate from other potential immunologically competent cells, and that at about the time of birth they migrate to the ultimate sites of immunological reactivity. Here, the thymocytes and their direct descendants, in the adult as well as in the neonate, are not themselves primarily immunologically reactive in the classic sense; however, in some way, as yet unknown they are capable of modifying or augmenting certain lines of lymphoid cells, i.e., by "hormonal" means, or by information transfer, thereby determining the extent, and increasing the specificity of their immunological reactivity.

Thus, one may regard the mammalian thymus as a recent immunological acquisition whose functions include discrimination of fine antigenic differences and the governance of reactivity of certain lines of less sophisticated lymphoid cells.

REFERENCES

1. Kalmutz, S. W., Nature 193, 851 (1962).
2. Miller, J. F. A. P., Lancet II, 748 (1961).
3. Martinez, C., Kersey, J., Papermaster, B. W., and Good, R. A.
Proc. Soc. Exp. Biol., N. Y. 109, 193 (1962).
4. Arnason, B. G., Jaukovic, B. D., and Waksman, B. H., Nature 194,
99, (1962).
5. Parrot, D. M. V., Transpl. Bull., 29, 102 (1962).
6. Delmasso, A. P., Martinez, C., Good, R. A., Proc. Soc. Exp. Biol.
and Med. 110, 205 (1962).
7. Papermaster, B. W., Delmasso, A. P., Martinez, C., Good, R. A.,
Proc. Soc. Exp. Biol. and Med. (1962).
8. Burnet, F. M., Proc. Royal Soc. Med. 55, 619 (1962).
9. Miller, J. F. A. P., Nature 195, 1318 (1962).
10. Congdon, C. C., Urso, I. S., Am. J. Path. 33, 749 (1957).
11. Urso, I. S., Rad. Res. 9, 197 (1958).
12. Barnes, D. W. H., Ilbery, P. L. T., and Loutit, J. F., Nature 181,
488 (1958).
13. Uphoff, D. E., J. Nat. Cancer Inst. 20, 625 (1958).
14. Tschetter, P. N., Githens, J. H., Moscovici, M. G., Blood 18, 182
(1961).
15. Doria, G., Goodman, J. W., Gengozian, N., and Congdon, C. C.,
J. Immun. 88, 20 (1962).
16. Cole, L. J., and Ellis, M. E., Science 128, 32 (1958).

DISTRIBUTION

Copies

NAVY

1-3	Chief, Bureau of Ships (Code 335)
4	Chief, Bureau of Ships (Code 320)
5-6	Chief, Bureau of Medicine and Surgery
7	Chief of Naval Operations (Op-07T)
8	Chief of Naval Research (Code 104)
9-11	Director, Naval Research Laboratory (Code 2021)
12	Office of Naval Research (Code 422)
13	Office of Naval Research (Code 441)
14-23	Office of Naval Research, FPO, New York
24-26	Naval Medical Research Institute
27	OIC, Radiation Exposure Evaluation Laboratory
28	Director, Aviation Medical Acceleration Laboratory
29	U.S. Naval Postgraduate School, Monterey
30	Commander, Naval Ordnance Laboratory, Silver Spring
31	Naval Missile Center (Code 5700)
32	U.S. Naval Hospital, San Diego
33	CO, Naval Medical Research Unit No. 2
34	CO, Naval Medical Field Research Laboratory, Camp Lejeune

ARMY

35	Chief of Research and Development (Atomic Division)
36	Chief of Research and Development (Life Science Division)
37	Deputy Chief of Staff for Military Operations (CBR)
38	Chief of Engineers (ENBMC-DE)
39	Chief of Engineers (ENG CW)
40	CG, Army Materiel Command (AMCRD-DE-NE)
41	CG, USA CBR Agency
42	President, Chemical Corps Board
43-45	CO, BW Laboratories
46	CO, Chemical Corps Training Command
47	Commandant, Chemical Corps Schools (Library)
48	CO, Chemical Research and Development Laboratories
49	Commander, Chemical Corps Nuclear Defense Laboratory
50	Hq., Army Environmental Hygiene Agency
51	CG, Aberdeen Proving Ground

52 CO, Army Medical Research Laboratory, Fort Knox
 53 Army Medical Research and Nutrition Laboratory (MEDEN-AD)
 54 CO, Army Medical Service Combat Development Agency
 55-56 Medical Field Service School, Fort Sam Houston (Stimson Lib.)
 57 Brooke Army Medical Center (Dept. Prev. Med.)
 58 Director, Surgical Research Unit, Fort Sam Houston
 59 Director, Walter Reed Army Medical Center
 60 Hq., Army Nuclear Medicine Research Detachment, Europe
 61 CG, Combat Developments Command (CDCMR-V)
 62 CG, Quartermaster Res. and Eng. Command
 63 Hq., Dugway Proving Ground
 64-66 The Surgeon General (MEDNE)
 67 Office of the Surgeon General (Combat Dev.)
 68 CG, Engineer Res. and Dev. Laboratory
 69 Director, Office of Special Weapons Development
 70 CG, Munitions Command
 71 CO, Frankford Arsenal
 72 CG, Army Missile Command

AIR FORCE

73 Assistant Chief of Staff, Intelligence (AFCIN-3B)
 74-79 CG, Aeronautical Systems Division (ASAPRD-NS)
 80 CO, Radiological Health Laboratory Division
 81 Director, USAF Project RAND
 82 Commandant, School of Aerospace Medicine, Brooks AFB
 83 CO, School of Aviation Medicine, Gunter AFB
 84 6571st Aeromedical Research Lab., Holloman AFB
 85 Radiobiological Laboratory
 86 Office of the Surgeon (SUP3.1), Strategic Air Command
 87 Office of the Surgeon General
 88 CG, Special Weapons Center, Kirtland AFB
 89 Director, Air University Library, Maxwell AFB
 90-91 Commander, Technical Training Wing, 3415th TTG
 92 Hq., Second Air Force, Barksdale AFB
 93 Commander, Electronic Systems Division (CRZT)

OTHER DOD ACTIVITIES

94-96 Chief, Defense Atomic Support Agency (Library)
 97 Commander, FC/DASA, Sandia Base (FCDV)
 98 Commander, FC/DASA, Sandia Base (FCTG5, Library)
 99 Commander, FC/DASA, Sandia Base (FCWT)
 100-101 Office of Civil Defense, Washington
 102-103 Civil Defense Unit, Army Library
 104 Armed Forces Institute of Pathology
 105-114 Armed Services Technical Information Agency
 115 Director, Armed Forces Radiobiology Research Institute

AEC ACTIVITIES AND OTHERS

116	Research Analysis Corporation
117	Life Science Officer, AEC, Washington
118	Director, Division of Biology and Medicine
119	NASA, Ames Research Center, Moffett Field
120	Naval Attache, Stockholm (for Commodore Troell)
121	Aerojet General, Azusa
122-126	Argonne Cancer Research Hospital
127-136	Argonne National Laboratory
137-138	Atomic Bomb Casualty Commission
139	AEC Scientific Representative, France
140	AEC Scientific Representative, Japan
141-143	Atomic Energy Commission, Washington
144-147	Atomic Energy of Canada, Limited
148-150	Atomics International
151-152	Battelle Memorial Institute
153	Borden Chemical Company
154-157	Brookhaven National Laboratory
158	Chicago Patent Group
159	Colorado State University
160	Columbia University (Rossi)
161	Committee on the Effects of Atomic Radiation
162-164	Defence Research Member
165-166	duPont Company, Aiken
167	duPont Company, Wilmington
168	Edgerton, Germeshausen and Grier, Inc., Goleta
169	Edgerton, Germeshausen and Grier, Inc., Las Vegas
170-171	General Dynamics, Fort Worth
172-173	General Electric Company, Cincinnati
174-181	General Electric Company, Richland
182	General Electric Company, St. Petersburg
183	General Scientific Corporation
184	Hughes Aircraft Company, Culver City
185	Iowa State University
186	Journal of Nuclear Medicine
187	Knolls Atomic Power Laboratory
188-189	Los Alamos Scientific Laboratory (Library)
190	Lovelace Foundation
191	Martin-Marietta Corporation
192	Massachusetts Institute of Technology
193-194	Midwestern Universities Research Association
195	Mound Laboratory
196	National Academy of Sciences
197-198	NASA, Scientific and Technical Information Facility
199	National Bureau of Standards (Taylor)
200	National Cancer Institute
201	National Lead Company of Ohio
202	National Library of Medicine
203	New Jersey State Department of Health

204 New York Operations Office
 205 New York University (Eisenbud)
 206 Office of Assistant General Counsel for Patents
 207-208 Phillips Petroleum Company
 209-212 Pratt and Whitney Aircraft Division
 213-214 Public Health Service, Washington
 215 Public Health Service, Las Vegas
 216 Public Health Service, Montgomery
 217 Sandia Corporation, Albuquerque
 218 Union Carbide Nuclear Company (ORGDG)
 219-223 Union Carbide Nuclear Company (ORNL)
 224 Union Carbide Nuclear Company (Paducah Plant)
 225 United Nuclear Corporation (NDA)
 226 U.S. Geological Survey, Denver
 227 U.S. Geological Survey, Menlo Park
 228 U.S. Geological Survey, Naval Weapons Plant
 229 U.S. Geological Survey, Washington
 230 U.S. Weather Bureau, Washington
 231 University of California, Davis
 232-234 University of California Lawrence Radiation Lab., Berkeley
 235-236 University of California Lawrence Radiation Lab., Livermore
 237 University of California, Los Angeles
 238 University of California, San Francisco
 239 University of Chicago Radiation Laboratory
 240 University of Hawaii
 241 University of Puerto Rico
 242 University of Rochester (Atomic Energy Project)
 243 University of Tennessee (UTA)
 244 University of Utah
 245 University of Washington (Donaldson)
 246 Wayne State University
 247 Westinghouse Electric Corporation (Rahilly)
 248 Westinghouse Electric Corporation (NASA)
 249 Western Reserve University (Friedell)
 250-274 Technical Information Extension, Oak Ridge

USNRDL

275-310 USNRDL, Technical Information Division

DISTRIBUTION DATE: 1 February 1963

<p>Naval Radiological Defense Laboratory USNRDL-TR-605</p> <p>MOUSE FETAL LIVER: A SOURCE OF IMMUNOLOGICALLY REACTIVE CELLS by M.L. Tyan and L.J. Cole 3 January 1963 17 p. table 16 refs.</p> <p>UNCLASSIFIED</p> <p>Male and female AHeJ mice were mated and the resulting pregnancies were surgically interrupted at various stages of gestation. The embryos were dissected and identical organs (liver, thymus, gut, placenta, and umbilicus) from each separate pregnancy were pooled and injected i.p. into sublethally irradiated (500 rad) F₁ hybrids (either (BALB/c x A))F₁ or (C57L x A))F₁). Death of the F₁ hybrids within 60 days was the criterion for the (over)</p>	<ol style="list-style-type: none"> 1. Liver. 2. Thymus. 3. Immunization. 4. Mammals. <ol style="list-style-type: none"> I. Tyan, M.L. II. Cole, L.J. III. Title. IV. MR005.08-5200. <p><u>UNCLASSIFIED</u></p>
---	--

<p>Naval Radiological Defense Laboratory USNRDL-TR-605</p> <p>MOUSE FETAL LIVER: A SOURCE OF IMMUNOLOGICALLY REACTIVE CELLS by M.L. Tyan and L.J. Cole 3 January 1963 17 p. table 16 refs.</p> <p>UNCLASSIFIED</p> <p>Male and female AHeJ mice were mated and the resulting pregnancies were surgically interrupted at various stages of gestation. The embryos were dissected and identical organs (liver, thymus, gut, placenta, and umbilicus) from each separate pregnancy were pooled and injected i.p. into sublethally irradiated (500 rad) F₁ hybrids (either (BALB/c x A))F₁ or (C57L x A))F₁). Death of the F₁ hybrids within 60 days was the criterion for the (over)</p>	<ol style="list-style-type: none"> 1. Liver. 2. Thymus. 3. Immunization. 4. Mammals. <ol style="list-style-type: none"> I. Tyan, M.L. II. Cole, L.J. III. Title. IV. MR005.08-5200. <p><u>UNCLASSIFIED</u></p>
---	--

<p>presence of immunologically competent cells in the inoculum. At the end of this period the survivors were sacrificed, their spleens and lymph nodes homogenized, and injected i.p. into sublethally irradiated F₁ hybrids of the other type. Again death within 60 days was the criterion for the presence of immunologically competent cells in the inoculum.</p> <p>One death occurred among the primary hosts (placenta 1/8). Immunologically competent cells (AHeJ) were detected in secondary recipients of 2nd trimester fetal liver (3/3), 2nd trimester thymus (3/3), and 3rd trimester fetal liver (1/12). No immunologically competent cells (AHeJ) were found in secondary recipients of 3rd trimester and newborn thymus (1/14). Deaths occurred among secondary hosts of fetal gut (3/5), umbilicus (1/3), and placenta (2/9). There were two deaths (2/29) among the secondary control mice. It is concluded that 2nd and 3rd trimester mouse fetal liver, 2nd trimester fetal thymus, and perhaps other fetal tissues contain potential immunologically competent cells. The theoretical significance of these data is discussed.</p>	<p><u>UNCLASSIFIED</u></p>
--	----------------------------

<p>presence of immunologically competent cells in the inoculum. At the end of this period the survivors were sacrificed, their spleens and lymph nodes homogenized, and injected i.p. into sublethally irradiated F₁ hybrids of the other type. Again death within 60 days was the criterion for the presence of immunologically competent cells in the inoculum.</p> <p>One death occurred among the primary hosts (placenta 1/8). Immunologically competent cells (AHeJ) were detected in secondary recipients of 2nd trimester fetal liver (3/3), 2nd trimester thymus (3/3), and 3rd trimester fetal liver (1/12). No immunologically competent cells (AHeJ) were found in secondary recipients of 3rd trimester and newborn thymus (1/14). Deaths occurred among secondary hosts of fetal gut (3/5), umbilicus (1/3), and placenta (2/9). There were two deaths (2/29) among the secondary control mice. It is concluded that 2nd and 3rd trimester mouse fetal liver, 2nd trimester fetal thymus, and perhaps other fetal tissues contain potential immunologically competent cells. The theoretical significance of these data is discussed.</p>	<p><u>UNCLASSIFIED</u></p>
--	----------------------------

<p>Naval Radiological Defense Laboratory USNRDL-TR-605</p> <p>MOUSE FETAL LIVER: A SOURCE OF IMMUNOLOGICALLY REACTIVE CELLS by M.L. Tyan and L.J. Cole 3 January 1963 17 p. table 16 refs.</p> <p>UNCLASSIFIED</p> <p>Male and female AHeJ mice were mated and the resulting pregnancies were surgically interrupted at various stages of gestation. The embryos were dissected and identical organs (liver, thymus, gut, placenta, and umbilicus) from each separate pregnancy were pooled and injected i.p. into sublethally irradiated (500 rad) F₁ hybrids (either (BALB/c x A))F₁ or (C57L x A))F₁). Death of the F₁ hybrids within 60 days was the criterion for the (over)</p>	<ol style="list-style-type: none"> 1. Liver. 2. Thymus. 3. Immunization. 4. Mammals. <ol style="list-style-type: none"> I. Tyan, M.L. II. Cole, L.J. III. Title. IV. MR005.08-5200. <p><u>UNCLASSIFIED</u></p>
---	--

<p>Naval Radiological Defense Laboratory USNRDL-TR-605</p> <p>MOUSE FETAL LIVER: A SOURCE OF IMMUNOLOGICALLY REACTIVE CELLS by M.L. Tyan and L.J. Cole 3 January 1963 17 p. table 16 refs.</p> <p>UNCLASSIFIED</p> <p>Male and female AHeJ mice were mated and the resulting pregnancies were surgically interrupted at various stages of gestation. The embryos were dissected and identical organs (liver, thymus, gut, placenta, and umbilicus) from each separate pregnancy were pooled and injected i.p. into sublethally irradiated (500 rad) F₁ hybrids (either (BALB/c x A))F₁ or (C57L x A))F₁). Death of the F₁ hybrids within 60 days was the criterion for the (over)</p>	<ol style="list-style-type: none"> 1. Liver. 2. Thymus. 3. Immunization. 4. Mammals. <ol style="list-style-type: none"> I. Tyan, M.L. II. Cole, L.J. III. Title. IV. MR005.08-5200. <p><u>UNCLASSIFIED</u></p>
---	--

<p>presence of immunologically competent cells in the inoculum. At the end of this period the survivors were sacrificed, their spleens and lymph nodes homogenized, and injected i.p. into sublethally irradiated F₁ hybrids of the other type. Again death within 60 days was the criterion for the presence of immunologically competent cells in the inoculum.</p> <p>One death occurred among the primary hosts (placenta 1/8). Immunologically competent cells (AHeJ) were detected in secondary recipients of 2nd trimester fetal liver (3/3), 2nd trimester thymus (3/3), and 3rd trimester fetal liver (1/12). No immunologically competent cells (AHeJ) were found in secondary recipients of 3rd trimester and newborn thymus (1/14). Deaths occurred among secondary hosts of fetal gut (3/5), umbilicus (1/3), and placenta (2/9). There were two deaths (2/29) among the secondary control mice. It is concluded that 2nd and 3rd trimester mouse fetal liver, 2nd trimester fetal thymus, and perhaps other fetal tissues contain potential immunologically competent cells. The theoretical significance of these data is discussed.</p>	<p><u>UNCLASSIFIED</u></p>
--	----------------------------

<p>presence of immunologically competent cells in the inoculum. At the end of this period the survivors were sacrificed, their spleens and lymph nodes homogenized, and injected i.p. into sublethally irradiated F₁ hybrids of the other type. Again death within 60 days was the criterion for the presence of immunologically competent cells in the inoculum.</p> <p>One death occurred among the primary hosts (placenta 1/8). Immunologically competent cells (AHeJ) were detected in secondary recipients of 2nd trimester fetal liver (3/3), 2nd trimester thymus (3/3), and 3rd trimester fetal liver (1/12). No immunologically competent cells (AHeJ) were found in secondary recipients of 3rd trimester and newborn thymus (1/14). Deaths occurred among secondary hosts of fetal gut (3/5), umbilicus (1/3), and placenta (2/9). There were two deaths (2/29) among the secondary control mice. It is concluded that 2nd and 3rd trimester mouse fetal liver, 2nd trimester fetal thymus, and perhaps other fetal tissues contain potential immunologically competent cells. The theoretical significance of these data is discussed.</p>	<p><u>UNCLASSIFIED</u></p>
--	----------------------------

<p>Naval Radiological Defense Laboratory USNRDL-TR-605</p> <p>MOUSE FETAL LIVER: A SOURCE OF IMMUNOLOGICALLY REACTIVE CELLS by M.L. Tyan and L.J. Cole 3 January 1963 17 p. table 16 refs.</p> <p>UNCLASSIFIED</p> <p>Male and female AHeJ mice were mated and the resulting pregnancies were surgically interrupted at various stages of gestation. The embryos were dissected and identical organs (liver, thymus, gut, placenta, and umbilicus) from each separate pregnancy were pooled and injected i.p. into sublethally irradiated (500 rad) F₁ hybrids (either (BALB/c x A))F₁ or (C57L x A))F₁). Death of the F₁ hybrids within 60 days was the criterion for the (over)</p>	<ol style="list-style-type: none"> 1. Liver. 2. Thymus. 3. Immunization. 4. Mammals. <ol style="list-style-type: none"> I. Tyan, M.L. II. Cole, L.J. III. Title. IV. MR005.08-5200. <p><u>UNCLASSIFIED</u></p>
---	--

<p>Naval Radiological Defense Laboratory USNRDL-TR-605</p> <p>MOUSE FETAL LIVER: A SOURCE OF IMMUNOLOGICALLY REACTIVE CELLS by M.L. Tyan and L.J. Cole 3 January 1963 17 p. table 16 refs.</p> <p>UNCLASSIFIED</p> <p>Male and female AHeJ mice were mated and the resulting pregnancies were surgically interrupted at various stages of gestation. The embryos were dissected and identical organs (liver, thymus, gut, placenta, and umbilicus) from each separate pregnancy were pooled and injected i.p. into sublethally irradiated (500 rad) F₁ hybrids (either (BALB/c x A))F₁ or (C57L x A))F₁). Death of the F₁ hybrids within 60 days was the criterion for the (over)</p>	<ol style="list-style-type: none"> 1. Liver. 2. Thymus. 3. Immunization. 4. Mammals. <ol style="list-style-type: none"> I. Tyan, M.L. II. Cole, L.J. III. Title. IV. MR005.08-5200. <p><u>UNCLASSIFIED</u></p>
---	--

<p>presence of immunologically competent cells in the inoculum. At the end of this period the survivors were sacrificed, their spleens and lymph nodes homogenized, and injected i.p. into sublethally irradiated F₁ hybrids of the other type. Again death within 60 days was the criterion for the presence of immunologically competent cells in the inoculum.</p> <p>One death occurred among the primary hosts (placenta 1/8). Immunologically competent cells (AHeJ) were detected in secondary recipients of 2nd trimester fetal liver (3/3), 2nd trimester thymus (3/3), and 3rd trimester fetal liver (1/12). No immunologically competent cells (AHeJ) were found in secondary recipients of 3rd trimester and newborn thymus (1/14). Deaths occurred among secondary hosts of fetal gut (3/5), umbilicus (1/3), and placenta (2/9). There were two deaths (2/29) among the secondary control mice. It is concluded that 2nd and 3rd trimester mouse fetal liver, 2nd trimester fetal thymus, and perhaps other fetal tissues contain potential immunologically competent cells. The theoretical significance of these data is discussed.</p>	<p><u>UNCLASSIFIED</u></p>
--	----------------------------

<p>presence of immunologically competent cells in the inoculum. At the end of this period the survivors were sacrificed, their spleens and lymph nodes homogenized, and injected i.p. into sublethally irradiated F₁ hybrids of the other type. Again death within 60 days was the criterion for the presence of immunologically competent cells in the inoculum.</p> <p>One death occurred among the primary hosts (placenta 1/8). Immunologically competent cells (AHeJ) were detected in secondary recipients of 2nd trimester fetal liver (3/3), 2nd trimester thymus (3/3), and 3rd trimester fetal liver (1/12). No immunologically competent cells (AHeJ) were found in secondary recipients of 3rd trimester and newborn thymus (1/14). Deaths occurred among secondary hosts of fetal gut (3/5), umbilicus (1/3), and placenta (2/9). There were two deaths (2/29) among the secondary control mice. It is concluded that 2nd and 3rd trimester mouse fetal liver, 2nd trimester fetal thymus, and perhaps other fetal tissues contain potential immunologically competent cells. The theoretical significance of these data is discussed.</p>	<p><u>UNCLASSIFIED</u></p>
--	----------------------------

<p>Naval Radiological Defense Laboratory USNRDL-TR-605</p> <p>MOUSE FETAL LIVER: A SOURCE OF IMMUNOLOGICALLY REACTIVE CELLS by M.L. Tyan and L.J. Cole 3 January 1963 17 p. table 16 refs. UNCLASSIFIED</p> <p>Male and female AHeJ mice were mated and the resulting pregnancies were surgically interrupted at various stages of gestation. The embryos were dissected and identical organs (liver, thymus, gut, placenta, and umbilicus) from each separate pregnancy were pooled and injected i.p. into sublethally irradiated (500 rad) F₁ hybrids (either (BALB/c x A))F₁ or ((C57L x A))F₁). Death of the F₁ hybrids within 60 days was the criterion for the (over)</p>	<ol style="list-style-type: none"> 1. Liver. 2. Thymus. 3. Immunization. 4. Mammals. I. Tyan, M.L. II. Cole, L.J. III. Title. IV. MR005.08-5200. <p style="text-align: center;"><u>UNCLASSIFIED</u></p>
<p>Naval Radiological Defense Laboratory USNRDL-TR-605</p> <p>MOUSE FETAL LIVER: A SOURCE OF IMMUNOLOGICALLY REACTIVE CELLS by M.L. Tyan and L.J. Cole 3 January 1963 17 p. table 16 refs. UNCLASSIFIED</p> <p>Male and female AHeJ mice were mated and the resulting pregnancies were surgically interrupted at various stages of gestation. The embryos were dissected and identical organs (liver, thymus, gut, placenta, and umbilicus) from each separate pregnancy were pooled and injected i.p. into sublethally irradiated (500 rad) F₁ hybrids (either (BALB/c x A))F₁ or ((C57L x A))F₁). Death of the F₁ hybrids within 60 days was the criterion for the (over)</p>	<ol style="list-style-type: none"> 1. Liver. 2. Thymus. 3. Immunization. 4. Mammals. I. Tyan, M.L. II. Cole, L.J. III. Title. IV. MR005.08-5200. <p style="text-align: center;"><u>UNCLASSIFIED</u></p>

presence of immunologically competent cells in the inoculum. At the end of this period the survivors were sacrificed, their spleens and lymph nodes homogenized, and injected i.p. into sublethally irradiated F₁ hybrids of the other type. Again death within 60 days was the criterion for the presence of immunologically competent cells in the inoculum.

One death occurred among the primary hosts (placenta 1/8). Immunologically competent cells (AHeJ) were detected in secondary recipients of 2nd trimester fetal liver (3/3), 2nd trimester thymus (3/3), and 3rd trimester fetal liver (11/12). No immunologically competent cells (AHeJ) were found in secondary recipients of 3rd trimester and newborn thymus (1/14). Deaths occurred among secondary hosts of fetal gut (3/5), umbilicus (1/3), and placenta (2/9). There were two deaths (2/29) among the secondary control mice. It is concluded that 2nd and 3rd trimester mouse fetal liver, 2nd trimester fetal thymus, and perhaps other fetal tissues contain potential immunologically competent cells. The theoretical significance of these data is discussed.

presence of immunologically competent cells in the inoculum. At the end of this period the survivors were sacrificed, their spleens and lymph nodes homogenized, and injected i.p. into sublethally irradiated F₁ hybrids of the other type. Again death within 60 days was the criterion for the presence of immunologically competent cells in the inoculum.

One death occurred among the primary hosts (placenta 1/8). Immunologically competent cells (AHeJ) were detected in secondary recipients of 2nd trimester fetal liver (3/3), 2nd trimester thymus (3/3), and 3rd trimester fetal liver (11/12). No immunologically competent cells (AHeJ) were found in secondary recipients of 3rd trimester and newborn thymus (1/14). Deaths occurred among secondary hosts of fetal gut (3/5), umbilicus (1/3), and placenta (2/9). There were two deaths (2/29) among the secondary control mice. It is concluded that 2nd and 3rd trimester mouse fetal liver, 2nd trimester fetal thymus, and perhaps other fetal tissues contain potential immunologically competent cells. The theoretical significance of these data is discussed.

UNCLASSIFIED

UNCLASSIFIED

Navy Department

**UNITED STATES NAVAL
RADIOLOGICAL DEFENSE
LABORATORY**

San Francisco 24, California

Official Business

Postage and Fees Paid
Navy Department

THIRD CLASS MAIL